

# Species Distribution Models in plant conservation science: a comprehensive review with a focus on Iran

Hooman Babanezhad<sup>1\*</sup>, Alireza Naqinezhad<sup>1,2</sup>

**Abstract** - This review article thoroughly examines the role of Species Distribution Models (SDMs) in plant conservation science, with a specific focus on applications within Iran. Commencing with an extensive methodological approach, involving an exhaustive search across reputable academic databases such as Scopus, Web of Science, and Google Scholar, the review synthesizes a comprehensive set of studies. It offers deep insights into SDM principles, challenges, and transformative applications. Addressing these challenges, the review explores contemporary data collection methods, including the use of remote sensing, drones, and citizen science, which enhance the precision and scope of SDMs. A detailed examination of various modeling algorithms and approaches, including MaxEnt, Random Forest, Bayesian models, and others, highlights their specific applications and contributions to plant conservation. The review also integrates climate change data and various scenarios into SDMs, showcasing case studies that illustrate SDMs' potential to predict shifts in plant distributions in response to changing climate conditions and overexploitation. Emphasizing the importance of spatial scale, the review discusses its critical impact on the accuracy of modeling and conservation planning. The article concludes by underlining the indispensable role of SDMs in advancing plant conservation efforts, offering tailored recommendations for researchers, policy-makers, and conservation practitioners.

**Key words:** climate change impacts, modeling algorithms, over-exploitation, plant conservation, spatial scale, Species Distribution Modeling.

**Riassunto** - *Species Distribution Model* nella conservazione della biodiversità vegetale: una rassegna completa con un focus sull'Iran.

Questo articolo di rassegna esamina a fondo il ruolo degli *Species Distribution Model* (SDM) nella conservazione della biodiversità vegetale, con un focus specifico sulle loro applicazioni in Iran. Partendo da un approccio metodologico approfondito, che ha comportato una ricerca esaustiva in rinomate banche dati accademiche come Scopus, Web of Science e Google Scholar, questa revisione sintetizza una serie completa di studi e offre una visione approfondita dei principi, delle sfide e delle applicazioni trasformative degli SDM. Per affrontare queste sfide, la rassegna

esplora i metodi di raccolta dati contemporanei, tra cui l'uso del telerilevamento, dei droni e della citizen science, che migliorano la precisione e la portata degli SDM. Un esame dettagliato di vari algoritmi e approcci di modellazione, tra cui MaxEnt, Random Forest, modelli bayesiani e altri, evidenzia le loro applicazioni specifiche e i contributi alla conservazione delle piante. La rassegna integra anche i dati sul cambiamento climatico e vari scenari negli SDM, mostrando casi di studio che illustrano il potenziale degli SDM nel prevedere i cambiamenti nella distribuzione delle piante in risposta al cambiamento delle condizioni climatiche e al sovrasfruttamento. Sottolineando l'importanza della scala spaziale, la rassegna discute il suo impatto critico sull'accuratezza della modellazione e sulla pianificazione della conservazione. L'articolo si conclude sottolineando il ruolo indispensabile degli SDM nel far progredire gli sforzi di conservazione della biodiversità vegetale, offrendo raccomandazioni su misura per i ricercatori, i responsabili politici e gli operatori della conservazione.

**Parole chiave:** algoritmi di modellazione, conservazione delle piante, impatto dei cambiamenti climatici, scala spaziale, sovrasfruttamento, *Species Distribution Modeling*.

## INTRODUCTION

Plants are the backbone of ecosystems, providing food, shelter, and oxygen to countless organisms, including humans (Pesce *et al.*, 2023). They contribute to nutrient cycling, soil formation, and climate regulation (Petsch *et al.*, 2023). The extinction or decline of plant species can have cascading effects on entire ecosystems, impacting wildlife, pollinators, and ecosystem services such as clean air and water (Plutino *et al.*, 2022). Recognizing this, plant conservation has gained increasing attention worldwide (Nyumba *et al.*, 2018; Palit *et al.*, 2022).

Plant conservation is a critical component of global biodiversity efforts, as plants form the foundation of terrestrial ecosystems and provide essential services to human societies (Mason *et al.*, 2022). However, the accelerating loss of plant species due to habitat destruction, climate change, invasive species, and other anthropogenic pressures has prompted an urgent need for effective conservation strategies. In this context, Species Distribution Modeling (SDM) has emerged as a powerful tool to inform and guide plant conservation efforts (Nicholson *et al.*, 2021).

SDM, also known as "Habitat Suitability Modeling" or "Ecological Niche Modeling" (Fragniere *et al.*, 2022), is a computational approach that leverages environmental and species occurrence data to predict the spatial distribution of species across landscapes. Initially developed in the field of biogeography, SDMs have evolved significantly over the past few decades, catalyzed by advancements in computa-

<sup>1</sup>Department of Plant Science, Faculty of Basic Sciences, University of Mazandaran, Babolsar, Iran

<sup>2</sup>Department of Environmental Sciences, College of Science and Engineering, University of Derby, UK.

\* Corresponding author: homan.babanejad@gmail.com

© 2025 Hooman Babanezhad, Alireza Naqinezhad

Received for publication: 21 May 2024

Accepted for publication: 27 September 2024

Online publication: 6 December 2024

tional techniques, data availability, and ecological understanding (Lippi *et al.*, 2019). In terms of practical applications, SDMs have been instrumental in guiding the conservation of numerous plant species and ecosystems. Examples include habitat restoration planning for rare and endangered plant species, invasive species management, and the design of protected areas to maximize plant diversity preservation (Bonebrake *et al.*, 2018; Rønsted *et al.*, 2023).

Early applications of SDMs in plant conservation predominantly focused on defining the habitat preferences of individual species and mapping their current distributions. Classic modeling algorithms such as MaxEnt and BIOCLIM paved the way for subsequent developments. Researchers employed these models to identify critical habitats, protect endangered species, and inform land-use planning (Kariyawasam *et al.*, 2019). Over time, the field witnessed a transition from single-species models to multispecies approaches, recognizing the ecological interdependencies among plants and their interactions with other organisms (Veselova & Gaziulusoy 2021). A significant leap forward has been the integration of high-resolution remote sensing data, which provides detailed information on vegetation cover, land use, and climate variables (Hamilton & Casey 2016). Furthermore, the incorporation of climate change projections into SDMs has become a vital aspect of contemporary plant conservation (Littlefield *et al.*, 2019). Spatial scale considerations are another noteworthy development in the field (Vilà-Cabrera *et al.*, 2019).

Climate change is a major driver of ecological shifts, impacting species distribution and biodiversity globally. Iran, with its diverse ecosystems and endemic species, faces significant challenges due to climate change. A study reveals significant altitudinal shifts in endemic species, emphasizing the importance of identifying high-diversity regions for targeted conservation efforts (Noori *et al.*, 2024). The warming Hyrcanian climate threatens the English yew in Northern Iran, with SDM projecting severe habitat loss by 2050 and 2070 (Alavi *et al.*, 2019). Management strategies, including assisted migration and genetic diversity conservation, are crucial for their survival (Erfanian *et al.*, 2021).

SDM predicts varying habitat suitability for Iranian *Prunus* species under different climate scenarios, emphasizing the need for tailored conservation and cultivation strategies to safeguard their genetic diversity (Zeraatkar & Khajoei Nasab, 2023). *Astragalus* genus distribution and diversity in Iran are identified as hotspot areas for conservation, emphasizing the need to address conservation gaps, particularly in regions not covered by global biodiversity hotspots (Maassoumi & Ashouri, 2022).

Climate change also impacts tree species in Hyrcanian forests, posing challenges for forest management and conservation (Niknaddaf *et al.*, 2023). SDM helps identify biodiversity hotspots, guiding conservation efforts in Hyrcanian Mountain Forests (Taleshi *et al.*, 2019; Ahmadi *et al.*, 2023; Mahmoodi *et al.*, 2023). Understanding tree species distributions aids in planning effective conservation strategies along elevation gradients. Lastly, SDM reveals the inadequacy of current protected areas in conserving endemic mountain flora in Iran, emphasizing the need for expansion and realignment of protected areas to mitigate future biodiversity loss (Limaki *et al.*, 2021).

Our review serves as a foundational step towards a more focused and comprehensive examination of SDMs in the context of plant conservation, particularly within Iran. While this review provides valuable insights into the broader field of SDMs and their practical applications, our ultimate goal is twofold. Firstly, we aim to conduct a more detailed review of SDM applications specifically tailored to Iran's unique biodiversity and conservation challenges. Secondly, we intend to identify the most effective SDM approaches for this specific context, ensuring that conservation strategies are optimally aligned with the distinct ecological and environmental characteristics of the region.

## METHODS

An extensive search was conducted across reputable academic databases, including Scopus, Web of Science, and Google Scholar. These databases were chosen for their comprehensive coverage of scientific literature. A search strategy was employed using a combination of relevant keywords and variations thereof. These search terms were used: "Species Distribution Modeling", "Habitat Suitability Modeling", "Ecological Niche Modeling", "Plant Conservation", "Plant Biodiversity", "Climate Change", "Overexploitations", "Land Use", "Remote Sensing", "Machine Learning", "MaxEnt", "Random Forest", "Gradient Boosting" and "Ensemble Modeling".

Articles that directly addressed the application of SDM to plant conservation or were closely related to the subject matter were considered. Inclusion was limited to articles published from the inception of SDM techniques to the latest available data up to September 2023. Only peer-reviewed articles from reputable academic journals, conference proceedings, or scholarly books were included in the review. Articles published in English were included due to accessibility for the intended readership. Relevant data were extracted from the selected articles, including study objectives, methodology, key findings, and implications for plant conservation.

Each selected article underwent a rigorous assessment to evaluate its quality and relevance. The aim was to ensure the inclusion of high-quality research and mitigate bias. Articles were evaluated for the rigor of their methodology and the clarity of their contributions to the field of plant conservation through SDMs. The findings and insights from the selected articles were synthesized and organized into thematic sections within the review article. Key advancements, trends, and emerging topics in SDMs for plant conservation were identified and presented.

## RESULTS AND DISCUSSION

To provide a comprehensive overview of the current state of SDMs in plant conservation, we reviewed a total of 64 articles. The selected studies span from the inception of SDM techniques in the early 2000s to the latest available data as of 2023. These articles were sourced from a range of academic databases, ensuring a broad representation of research within the field. The geographic context of the studies is diverse, covering 21 countries. Notably, the highest number of studies originated from the USA (15 studies), followed by China (8 studies) and Iran (6 studies), as plotted

in Fig. 1. This geographic distribution underscores the global relevance of SDMs while also highlighting regions where these techniques have been prominently applied.

To provide a clear understanding of the temporal distribution of research, we analyzed the number of studies published over time. The data reveals a significant increase in publications, with a peak in 2017 when 10 studies were published (Fig. 2). This trend indicates growing interest and advancements in the application of SDMs in plant conservation.

### Advancements in data collection

Data collection is a fundamental component of SDMs, and recent advancements in technology and methodologies have greatly improved the quality and quantity of data available for modeling plant species distributions.

Herbarium specimens undergo a remarkable transformation as they journey into the digital realm, becoming accessible resources through platforms like GBIF and SEINet. These repositories burgeon with images and invaluable data, spawning a wave of opportunities (Marsico *et al.*, 2020; Heberling, 2022). Rare plant specimens, once confined to cabinets, now fuel the development of habitat suitability models and prognosticate range shifts (Molano-Flores *et al.*, 2023). Extracted occurrence data from these specimens emerge as linchpins for species regression analyses and the assessment of conservation statuses (Besnard *et al.*, 2018). Portals such as iDigBio, GBIF, and others burgeon with this wealth of information, serving as vital reservoirs that plug gaps in the distribution of elusive plants (Saran *et al.*, 2022). In the do-

main of SDMs, these repositories serve as pivotal data fountains, frequently tapped for species occurrence data modeling (Ribeiro *et al.*, 2022). Moreover, herbaria-based localities prove invaluable in strategic species assessments, acting as substitutes or supplements in modeling species distributions. They guide field surveys, unveil new populations (de Queiroz *et al.*, 2012; McCune, 2016), foretell future range changes (Erfanian *et al.*, 2021; Shay *et al.*, 2021), dissect climate change impacts (Abbott *et al.*, 2017; Dangremond *et al.*, 2022), and play a crucial role in shaping conservation strategies for rare species (Qazi *et al.*, 2022).

Remote sensing technologies, including satellite and aerial imagery, have provided a wealth of high-resolution data on vegetation cover, land use, and climate variables. This spatially explicit information is invaluable for characterizing the environment (Pricope *et al.*, 2019). Unmanned aerial vehicles equipped with sensors and cameras enable researchers to collect fine-scale, real-time data on plant communities and habitat characteristics. These platforms are particularly useful for monitoring and assessing plant populations in remote or challenging terrain (Castellanos-Galindo *et al.*, 2019). Citizen science initiatives and crowdsourced data collection platforms have expanded the spatial and temporal coverage of species observations. These efforts engage the public in data collection, providing valuable insights into plant distributions and phenology (Giovos *et al.*, 2019).

Environmental DNA (eDNA) techniques involve the collection and analysis of genetic material shed by organisms into their environment. In plant conservation, eDNA can be used to detect the presence of rare or elusive plant




Fig. 1 – Distribution of studies by country. / Distribuzione degli studi per nazione.

species in aquatic or terrestrial habitats (Johnson *et al.*, 2023). Camera traps and acoustic recording devices are increasingly used to capture data on plant-associated wildlife, which can indirectly inform plant distribution models by providing insights into herbivory, seed dispersal, and pollination (Caravaggi *et al.*, 2020). Field-deployed sensors can capture real-time data on environmental conditions, including soil moisture, temperature, and light levels. These sensors contribute to more accurate and continuous monitoring of plant habitats (Rühm *et al.*, 2023).

### Modeling algorithms and approaches

SDM relies on a variety of modeling algorithms and approaches, each with its strengths and limitations. Advances in computational methods have provided ecologists and conservationists with a diverse toolkit for understanding and predicting plant species distributions.

MaxEnt is a widely used algorithm known for its ability to handle presence-only data effectively. It estimates a probability distribution by maximizing entropy subject to constraints defined by environmental variables (Gao *et al.*, 2021). Random Forest is an ensemble machine learning method that combines multiple decision trees to make predictions. It is robust, handles complex interactions, and can accommodate both presence-absence and presence-only data (Gao & Zhou, 2020). Gradient boosting algorithms, such as XGBoost and LightGBM, have gained popularity for their predictive accuracy and ability to handle large da-

taset. They are used in SDMs to model complex relationships between species and environmental variables (Chen *et al.*, 2020).

GLMs provide a statistical framework for modeling species distributions and are appreciated for their interpretability. They are particularly useful when simpler models are preferred or when the relationships between variables are well-understood (Lee, 2021). Deep learning methods, including neural networks and convolutional neural networks, are increasingly applied to SDMs, especially for image-based data and complex spatial relationships (Brunsdon, 2020). Bayesian modeling approaches, such as hierarchical models and Bayesian networks, are used in SDMs to incorporate uncertainty and account for complex spatial dependencies (Oyster *et al.*, 2018). Ensemble modeling combines predictions from multiple algorithms to improve accuracy and robustness. It is a common practice in SDMs to mitigate the limitations of individual algorithms (Tanner *et al.*, 2017). Functional trait-based modeling focuses on the traits of plant species and how they relate to environmental conditions. This approach can provide mechanistic insights into plant distributions (Classen *et al.*, 2017).

Incorporating demographic data into SDMs, especially under climate change scenarios, offers a more refined understanding of species' potential distributions and extinction risks. Demographic factors such as growth rates, survival, and reproductive success play crucial roles in species resili-




Fig. 2 – Number of studies employing Species Distribution Models published from 2000 to 2023. / Numero degli studi che hanno utilizzato gli Species Distribution Models pubblicati dal 2000 al 2023.



ience to changing environments demonstrated this by integrating demographic data into SDMs for high-mountain plants, which allowed for better predictions of species' future distributions and informed more targeted conservation strategies (Dullinger *et al.*, 2012).

### **Incorporating climate change: integration into plant Species Distribution Models**

Incorporating climate change data and scenarios into plant SDMs is critical for understanding how shifting environmental conditions may impact plant species distributions and, consequently, for effective conservation planning. Recent advancements in climate modeling and ecological modeling techniques have allowed for more accurate assessments of these potential changes.

Integrating climate data into SDMs involves using historical climate records and future climate projections to assess how changing temperature and precipitation patterns will influence plant distributions (Araújo *et al.*, 2019). Researchers often use various emission scenarios, such as those outlined by the Intergovernmental Panel on Climate Change, to model potential climate futures and their impact on plant species (Masson-Delmotte *et al.*, 2021). In a study assessing alpine plant species in the European Alps, researchers used SDMs and climate change scenarios to predict upward shifts in species' ranges due to warming temperatures (Harvey *et al.*, 2017). A study on oak tree species in the United States utilized SDMs and climate projections to predict significant shifts in oak species ranges over the next century, highlighting potential conservation challenges (Iverson *et al.*, 2019).

Researchers have used climate data to identify vulnerable plant species and predict how changes in temperature and precipitation might affect their distributions. This information is vital for conservation planning (Beaumont *et al.*, 2016). Some studies go beyond single-species modeling to investigate community-level responses to climate change, considering interactions among multiple plant species (Ovaskainen *et al.*, 2016). Research on high-altitude plants in the southern Alps of New Zealand used SDMs to project range contractions due to climate warming, emphasizing the need for conservation action (McPherson, 2022).

### **Spatial scale and plant conservation**

Spatial scale plays a crucial role in SDM, especially in the context of plant conservation. The choice of scale can significantly influence the accuracy of predictions and the effectiveness of conservation planning.

Spatial scale refers to the size or extent of the geographic area over which data are collected and analyzed in SDMs. It is a fundamental consideration because it determines the level of detail and precision in modeling plant species distributions (Wang *et al.*, 2017). Fine-scale models focus on small geographic areas and provide detailed information about local habitat suitability. These models are valuable for site-specific conservation decisions and understanding the microhabitat preferences of plant species. Broad-scale models cover larger geographic regions and are useful for identifying broad distribution patterns and regional conservation priorities. They are often employed for landscape-level planning (Danino *et al.*, 2016). Different ecological processes operate at differ-

ent scales. For example, seed dispersal may be influenced by fine-scale landscape features, while climate change impacts may manifest at broader scales. Choosing the appropriate scale aligns modeling with the relevant ecological processes (Rocchini *et al.*, 2017). The mismatch between the scale of data collection and the scale of ecological processes can lead to ecological fallacies. For instance, making local conservation decisions based on broad-scale models can result in ineffective strategies (Simião-Ferreira *et al.*, 2018).

To address the scale issue, researchers often employ multi-scale modeling approaches. These approaches integrate information from different spatial scales to provide a comprehensive understanding of species distributions (Sanguet *et al.*, 2022). In plant conservation, spatial scale considerations are vital when prioritizing protected areas, designing corridors for habitat connectivity, and assessing the impacts of land-use changes (Liang & Song, 2022). Climate change modeling often requires dynamic scaling, considering temporal scales alongside spatial scales, to project how plant species distributions may shift over time (Graham *et al.*, 2019).

The effectiveness of SDMs varies significantly with the spatial scale of application. Fine-scale models, such as those applied in microhabitat studies, can capture detailed species responses to local environmental gradients and microclimatic variations. For example, Steinbauer *et al.* (2018) used fine-scale SDMs to analyze the impacts of climate change on plant species in the European Alps, highlighting how these models can identify microrefugia where species might survive despite broader climatic shifts. On the other hand, broad-scale models are more suitable for understanding regional shifts in species distributions and are often used in large-scale conservation planning.

### **Conservation applications of Species Distribution Models in plant**

SDMs are powerful tools in plant conservation, providing critical insights into habitat suitability, potential distribution shifts, and identifying priority areas for conservation efforts. This section delves into various real-world applications of SDMs, illustrating their utility in conservation planning and management.

Remote sensing data can significantly enhance SDMs by supplying comprehensive environmental data such as land cover, vegetation indices, and climate variables. This data provides broad-scale context, helping to infer habitat conditions and landscape features crucial for modeling species distributions. However, remote sensing often lacks species-specific detail, which may limit the accuracy of ecological niche modeling, particularly for species with specialized habitat requirements. To address these limitations, combining remote sensing with ground-truthing and high-resolution ecological data is essential, allowing for more precise modeling of species distributions.

SDMs are also valuable in managing invasive species by predicting potential distributions based on historical data and assessing habitat permeability. For instance, Van Nuland *et al.* (2016) utilized SDMs to predict the spread of invasive plants in the Hawaiian Islands, aiding in the development of strategies to mitigate their impact on vulnerable native habitats.

## Case Studies

### *Orchid conservation in China*

In their study, Li *et al.* (2022) used MaxEnt, a machine learning SDM, to assess habitat suitability for endangered orchid species such as *Cymbidium tortisepalum* Fukuy and *Paphiopedilum dianthum* Tang & F.T.Wang. By integrating environmental variables such as temperature, precipitation, and soil type, the model identified key conservation areas with high habitat suitability. This detailed approach helps prioritize sites for in-situ conservation and potential reintroduction programs, aligning closely with the theoretical frameworks discussed earlier.

### *Climate change impact in the European Alps*

Steinbauer *et al.* (2018) employed ensemble modeling, combining outputs from multiple SDMs to predict potential range shifts and habitat loss for rare alpine plants under various climate change scenarios. This method, integrating fine-scale climate and topographic data, allowed for a comprehensive assessment of future habitat suitability and the identification of potential microrefugia, providing actionable insights for conservation planning.

### *Restoration in Australia's Murray-Darling Basin*

In this case, Gawne *et al.* (2020) used SDMs based on Bioclim and other correlative models to guide the selection of planting sites for native species. SDMs incorporated data on soil moisture, elevation, and historical plant distributions to identify areas most suitable for restoration efforts. This approach enhanced habitat restoration by ensuring that plantings were conducted in areas with optimal conditions for survival and growth.

### *Conservation in the Azores Archipelago*

Rocha-Ortega *et al.* (2020) applied MaxEnt models to prioritize conservation actions for rare endemic plants in the Azores. By analyzing species distribution against climate variables and land use data, the study identified critical areas for establishing new protected zones, ensuring the conservation of these unique species in the face of environmental change.

### *Enhancing habitat connectivity in fragmented landscapes*

Liu *et al.* (2018) utilized SDMs to assess and enhance habitat connectivity for plant species in fragmented landscapes. By modeling potential movement corridors and identifying habitat patches critical for maintaining genetic flow, this study provided a foundation for conservation actions aimed at reducing species isolation and promoting long-term survival.

## Species Distribution Models in Iran

SDMs have been instrumental in understanding the distribution and conservation needs of various plant species in Iran. The following case studies illustrate the effectiveness of SDMs in guiding conservation strategies and management actions across different regions and ecosystems in Iran.

### *Nepeta crispa in western Iran*

The study on *Nepeta crispa* Willd. demonstrated the effectiveness of SDMs in identifying current habitat suit-

ability and forecasting future range shifts due to climate change. The model indicated that elevation, temperature, geology, and precipitation are critical factors influencing the distribution of *N. crispa*. As a result, it was suggested that conservation efforts should prioritize areas above 2000 m a.s.l., where suitable habitats are likely to persist under changing climatic conditions (Mahmoodi *et al.*, 2022).

### *Brant's oak and climate change*

Research on *Quercus brantii* Lindl. in western Iran utilized ensemble modeling to assess the potential impacts of climate change on the species' distribution. The results revealed a significant vulnerability of *Q. brantii* to climate change, projecting a substantial loss of suitable habitats by 2070 under the most pessimistic scenario. This emphasizes the urgency for proactive conservation measures, such as protecting climate refugia and enhancing habitat connectivity, to mitigate the anticipated range contractions (Safaei *et al.*, 2021).

### *Crop wild relatives and Aegilops species*

The study on crop wild relatives, particularly focusing on goatgrasses *Aegilops* L. species, employed MaxEnt modeling to predict range changes under various climatic scenarios. The findings indicated mixed responses, with some species experiencing range contractions and other expansions. This variability underscores the need for dynamic conservation planning that can adapt to the changing distributions of these valuable genetic resources, ensuring their preservation and potential utility for future crop breeding (Hosseini *et al.*, 2022).

### *Hyrcanian temperate forests*

A study in the Hyrcanian temperate forests highlighted the importance of understorey plant species for biodiversity conservation. SDMs were used to predict the occurrence of these species based on climate, soil, and canopy cover variables. The results showed that while climate variables are pivotal in determining species distributions, changes in canopy cover due to forest management could either mitigate or exacerbate the effects of climate change on different understorey plant groups. This points to the need for tailored forest management practices that consider the specific ecological requirements of shade-adapted and sun-adapted species (Naqinezhad *et al.*, 2022).

### *Artemisia sieberi in central Iran*

Research on *Artemisia sieberi* Besser in central Iran compared the effectiveness of SDMs using bioclimatic variables derived from remote sensing data versus traditional instrumental records. The study found that models incorporating remote sensing data significantly outperformed those based solely on instrumental records, highlighting the value of up-to-date, high-resolution data for enhancing model accuracy. This demonstrates the potential of remote sensing technologies to improve conservation planning by providing reliable predictions for habitat management (Amiri *et al.*, 2020).

### *Brant's oak distribution across Iran*

The study on the distribution of Brant's oak *Quercus brantii* Lindl. across different geographical extents used var-

ious SDMs, including Random Forest, Generalized Linear Model, and Maximum Entropy, to project future habitat suitability under climate change scenarios. The results indicated a likely decline in potential habitats for Brant's oak across all examined spatial scales, reinforcing the need for scale-specific conservation strategies that address both local and regional challenges in habitat preservation (Mirhashemi *et al.*, 2023).

Overall, these case studies underscore the critical role of SDMs in informing conservation strategies in Iran. By providing precise predictions of species distributions and identifying priority areas for conservation, SDMs offer essential guidance for developing effective management plans to protect biodiversity in diverse ecosystems.

### Challenges and limitations of Species Distribution Models in plant conservation

High-quality data on plant occurrences and environmental variables are crucial for accurate SDMs. However, such data may be scarce, biased, or incomplete, especially in remote or understudied regions (Liu *et al.*, 2019). The choice of spatial scale in SDMs can be challenging, and mismatches between the scale of the model and the scale of ecological processes can lead to inaccurate predictions (Shaikh *et al.*, 2021). In many SDM analyses, one could assume that species are in equilibrium with their environment, but this assumption may not hold true, especially in rapidly changing environments due to factors like climate change or habitat degradation (Adhikari *et al.*, 2019). Incorporating climate change scenarios into SDMs introduces uncertainty, as future climate conditions are unpredictable. This can affect the accuracy of predictions related to plant distribution shifts (Yu *et al.*, 2019).

SDMs often do not account for biotic interactions, such as competition or mutualism, which can significantly influence plant distributions (Wen *et al.*, 2016). Overfitting occurs when models are overly complex and fit noise in the data, leading to poor generalization to new locations or conditions (Lee-Yaw *et al.*, 2022). Models built in one geographic region may not perform well in others due to differences in environmental conditions, leading to challenges in transferring knowledge across regions (Sillero & Barbosa, 2021). SDMs often assume species' distributions are static, but evolutionary processes, such as adaptation and migration, can influence plant distribution patterns over time (Escobar & Craft, 2016).

The potential applicability of SDMs in identifying conservation gaps within countries is a significant aspect worth highlighting. SDMs can serve as valuable tools for identifying areas that are underrepresented or not adequately covered within conservation efforts (Cha *et al.*, 2021; Zhong *et al.*, 2021). By utilizing SDMs, researchers and conservationists can pinpoint regions where species richness is high, but conservation efforts are low or insufficient. These models can help prioritize areas for conservation interventions, directing resources toward areas that are ecologically significant yet overlooked in conservation planning.

In the context of Iran, for instance, employing SDMs can aid in identifying specific regions that harbor unique or endangered plant species but are currently not adequately protected or considered in conservation strategies. This knowledge is instrumental in guiding policymaking and re-

source allocation to fill these conservation gaps and ensure the preservation of the country's rich plant diversity. Highlighting the role of SDMs in pinpointing conservation gaps within countries like Iran adds depth to the discussion by showcasing the practical utility of these models beyond predictive purposes, emphasizing their crucial role in guiding targeted and effective conservation actions.

### CONCLUSIONS

SDMs indeed possess a versatile applicability beyond plant conservation, extending their utility across diverse organismic groups and ecosystems. Their adaptability allows for the exploration and understanding of species distributions and environmental relationships across various taxa, including animals, microbes, and even entire ecological communities. Whether studying the habitat preferences of mammals, the range shifts of birds due to climate change, or the distributions of disease vectors, SDMs serve as powerful tools for understanding and predicting species' responses to environmental changes. Moreover, these models find applications in conservation biology, epidemiology, invasive species management, and ecosystem restoration, among others. For instance, in conservation, SDMs aid in identifying critical habitats for endangered species and help design protected areas. In epidemiology, they assist in mapping disease risk and understanding disease transmission dynamics. Their versatility in predicting species distributions and understanding ecological relationships makes SDMs invaluable across multiple fields of study and conservation efforts.

In conclusion, SDMs have proven to be invaluable tools for advancing plant conservation efforts by offering a data-driven approach to comprehending, projecting, and mitigating the effects of environmental changes on plant species. Recognizing the pivotal role of SDMs, recommendations for various stakeholders emerge. Researchers are encouraged to persist in refining and enhancing SDM techniques while tackling data quality issues and embracing new data sources. Further exploration into integrating biotic interactions and evolutionary dynamics into SDMs is advocated to augment predictive accuracy. Policymakers are urged to acknowledge the value of SDMs in conservation planning, earmarking resources for data collection, modeling, and monitoring endeavors, with an emphasis on incorporating SDM-based predictions into policy and land-use planning to ensure the sustainability of plant conservation efforts. Conservation practitioners are encouraged to harness the potential of SDMs to steer on-the-ground conservation actions, encompassing habitat restoration, invasive species management, and the design of protected areas. Additionally, fostering collaborations with researchers is pivotal to ensuring the seamless integration of the latest SDM advancements into conservation strategies, culminating in a holistic approach to safeguarding plant biodiversity.

### REFERENCES

- Abbott R.E., Doak D.F. & DeMarche M.L., 2017 – Portfolio effects, climate change, and the persistence of small populations: analyses on the rare plant *Saussurea Weberi*. *Ecology*, 98 (4): 1071-1081.



- Adhikari A., Mainali K.P., Rangwala I. & Hansen A.J., 2019 – Various measures of potential evapotranspiration have species-specific impact on species distribution models. *Ecological Modelling*, 414: 108836.
- Ahmadi K., Mahmoodi S., Pal S.C., Saha A., Chowdhuri I., Kolyaie S., Linh N.T.T. & Kumar L., 2023 – Modeling tree species richness patterns and their environmental drivers across Hyrcanian mountain forests. *Ecological Informatics*, 77: 102226.
- Alavi S.J., Ahmadi K., Hosseini S.M., Tabari M. & Nouri Z., 2019 – The response of English yew (*Taxus baccata* L.) to climate change in the Caspian Hyrcanian mixed forest ecoregion. *Regional Environmental Change*, 19: 1495-1506.
- Amiri M., Tarkesh M., Jafari R. & Jetschke G., 2020 – Bioclimatic variables from precipitation and temperature records vs. remote sensing-based bioclimatic variables: which side can perform better in species distribution modeling? *Ecological Informatics*, 57: 101060.
- Araújo M.B., Anderson R.P., Márcia Barbosa A., Beale C.M., Dormann C.F., Early R., Garcia R.A., Guisan A., Maiorano L., Naimi B., O'Hara R.B., Zimmermann N.E. & Rahbek C., 2019 – Standards for distribution models in biodiversity assessments. *Science Advances*, 5 (1): eaat4858.
- Beaumont L.J., Graham E., Duursma D.E., Wilson P.D., Cabrelli A., Baumgartner J.B., Hallgren W., Esperón-Rodríguez M., Nipperess D.A., Warren D.L., Laffan S.W. & VanDerWal J., 2016 – Which species distribution models are more (or less) likely to project broad-scale, climate-induced shifts in species ranges? *Ecological Modelling*, 342: 135-146.
- Besnard G., Gaudeul M., Lavergne S., Muller S., Rouhan G., Sukhorukov A.P., Vanderpoorten A. & Jabbour F., 2018 – Herbarium-based science in the twenty-first century. *Botany Letters*, 165 (3-4): 323-327.
- Bonebrake T.C., Brown C.J., Bell J.D., Blanchard J.L., Chauvenet A., Champion C., Chen I.C., Clark T.D., Colwell R.K., Danielsen F., Dell A.I., Donelson J.M., Evengård B., Ferrier S., Frusher S., Garcia R.A., Griffis R.B., Hobday A.J., Jarzyna M.A., Lee E., Lenoir J., Linnetved H., Martin V.Y., McCormack P.C., McDonald J., McDonald-Madden E., Mitchell N., Mustonen T., Pandolfi J.M., Pettorelli N., Possingham H., Pulsifer P., Reynolds M., Scheffers B.R., Sorte C.J.B., Strugnell J.M., Tuanmu M.N., Twiname S., Vergés A., Villanueva C., Wapstra E., Wernberg T. & Pecl G.T., 2018 – Managing consequences of climate-driven species redistribution requires integration of ecology, conservation and social science. *Biological Reviews*, 93: 284-305.
- Brunsdon C., 2020 – Robin Lovelace, Jakub Nowosad and Jannes Muenchow. Geocomputation with R. SAGE Publications Sage, London.
- Caravaggi A., Burton A.C., Clark D.A., Fisher J.T., Grass A., Green S., Hobaiter C., Hofmeester T.R., Kalan A.K., Rabaiotti D. & Rivet D., 2020 – A review of factors to consider when using camera traps to study animal behavior to inform wildlife ecology and conservation. *Conservation Science and Practice*, 2 (8): e239.
- Castellanos-Galindo G.A., Casella E., Mejía-Rentería J.C. & Rovere A., 2019 – Habitat mapping of remote coasts: evaluating the usefulness of lightweight unmanned aerial vehicles for conservation and monitoring. *Biological Conservation*, 239: 108282.
- Cha Y., Shin J., Go B., Lee D.S., Kim Y., Kim T. & Park Y. S., 2021 – An interpretable machine learning method for supporting ecosystem management: application to species distribution models of freshwater macroinvertebrates. *Journal of Environmental Management*, 291: 112719.
- Chen R.C., Caraka R.E., Arnita N.E.G., Pomalingo S., Rachman A., Toharudin T., Tai S.K. & Pardamean B., 2020 – An end to end of scalable tree boosting system. *Sylwan*, 164: 140-150.
- Classen A., Steffan-Dewenter I., Kindeketa W.J. & Peters M.K., 2017 – Integrating intraspecific variation in community ecology unifies theories on body size shifts along climatic gradients. *Functional Ecology*, 31 (3): 768-777.
- Dangremond E.M., Hill C.H., Louaibi S. & Munoz I., 2022 – Phenological responsiveness and fecundity decline near the southern range limit of *Trientalis borealis* (Primulaceae). *Plant Ecology*, 223: 41-52.
- Danino M., Shnerb N.M., Azale S., Kunin W.E. & Kessler D.A., 2016 – The effect of environmental stochasticity on species richness in neutral communities. *Journal of Theoretical Biology*, 409: 155-164.
- De Queiroz T.F., Baughman C., Baughman O., Gara M. & Williams N., 2012 – Species distribution modeling for conservation of rare, edaphic endemic plants in White River Valley, Nevada. *Natural Areas Journal*, 32 (2): 149-158. <https://doi.org/10.3375/043.032.0203>
- Dullinger S., Gattlinger A., Thuiller W., Moser D., Zimmermann N. E., Guisan A., Willner W., Plutzar C., Leitner M., Hülber K., et al., 2012 – Extinction debt of high-mountain plants under twenty-first-century climate change. *Nature Climate Change*, 2 (8): 619-622.
- Erfanian M.B., Sagharyan M., Memariani F. & Ejtehadi H., 2021 – Predicting range shifts of three endangered endemic plants of the Khorassan-Kopet Dagh floristic province under global change. *Scientific Reports*, 11: 9159.
- Escobar L.E. & Craft M.E., 2016 – Advances and limitations of disease biogeography using ecological niche modeling. *Frontiers in Microbiology*, 7: 1174.
- Fragniere Y., Gremaud J., Pesenti E., Bétrisey S., Petitpierre B., Guisan A. & Kozłowski G., 2022 – Mapping habitats sensitive to overgrazing in the Swiss northern Alps using habitat suitability modeling. *Biological Conservation*, 274: 109742.
- Gao T., Xu Q., Liu Y., Zhao J. & Shi J., 2021 – Predicting the potential geographic distribution of *Sirex nitobei* in China under climate change using maximum entropy model. *Forests*, 12 (2): 151.
- Gao W. & Zhou Z., 2020 – Towards convergence rate analysis of random forests for classification. *Advances in neural information processing systems*, 33: 9300-9311.
- Gawne B., Hale J., Stewardson M.J., Webb J.A., Ryder D.S., Brooks S.S., Campbell C.J., Capon S.J., Everingham P., Grace M.R., et al., 2020 – Monitoring of environmental flow outcomes in a large river basin: the commonwealth environmental water holder's long-term intervention in the Murray-Darling basin, Australia. *River Research and Applications*, 36 (4): 630-644.
- Giovos I., Kleitou P., Poursanidis D., Batjakas I., Bernardi G., Crocetta F., Doumpas N., Kalogirou S., Kampouris T.E., Keramidas I., Lange-neck J., Maximidi M., Mitsou E., Stoilas V.O., Tiralongo F., Romanidis-Kyriakidis G., Xentidis N.J., Zenetos A. & Katsanevakis S., 2019 – Citizen-science for monitoring marine invasions and stimulating public engagement: a case project from the Eastern Mediterranean. *Biological Invasions*, 21: 3707-3721.
- Graham V., Baumgartner J.B., Beaumont L.J., Esperón-Rodríguez M. & Grech A., 2019 – Prioritizing the protection of climate refugia: designing a climate-ready protected area network. *Journal of Environmental Planning and Management*, 62 (14): 2588-2606.
- Hamilton S.E. & Casey D., 2016 – Creation of a high spatio-temporal resolution global database of continuous mangrove forest cover for the 21st century (Cgmfc-21). *Global Ecology and Biogeography*, 25 (6): 729-738.
- Harvey E., Gounand I., Ward C.L. & Altermatt F., 2017 – Bridging ecology and conservation: from ecological networks to ecosystem function. *Journal of Applied Ecology*, 54 (2): 371-379.
- Heberling J.M., 2022 – Herbaria as big data sources of plant traits. *International Journal of Plant Sciences*, 183 (2): 87-118.
- Hosseini N., Mehrabian A. & Mostafavi H., 2022 – Modeling climate change effects on spatial distribution of wild *Aegilops* L. (Poaceae) toward food security management and biodiversity conservation in Iran. *Integrated Environmental Assessment and Management*, 18 (3): 697-708.
- Iverson L.R., Prasad A.M., Peters M.P. & Matthews S.N., 2019 – Facilitating adaptive forest management under climate change: a spatially specific synthesis of 125 species for habitat changes and assisted migration over the eastern United States. *Forests*, 10 (11): 989.
- Johnson M.D., Freeland J.R., Parducci L., Evans D.M., Meyer R.S., Molano-Flores B. & Davis M.A., 2023 – Environmental DNA as an emerging tool in botanical research. *American Journal of Botany*, 110 (2): e16120.



- Kariyawasam C.S., Kumar L. & Ratnayake S.S., 2019 – Invasive plants distribution modeling: a tool for tropical biodiversity conservation with special reference to Sri Lanka. *Tropical Conservation Science*, 12 (6): 194008291986426.
- Lee-Yaw J.A., McCune J.L., Pironon S. & Sheth S.N., 2022 – Species distribution models rarely predict the biology of real populations. *Ecography*, 2022 (6): e05877.
- Lee Y., 2021 – Beyond multiple linear regression: applied generalized linear models and multilevel models in R. *Taylor & Francis*.
- Li W., Zhao Q., Guo M., Lu C., Huang F., Wang Z. & Niu J., 2022 – Predicting the potential distribution of the endangered plant *Cremastra Appendiculata* (orchidaceae) in China under multiple climate Change Scenarios. *Forests*, 13 (9): 1504.
- Liang Y. & Song W., 2022 – Integrating potential ecosystem services losses into ecological risk assessment of land use changes: a case study on the Qinghai-Tibet plateau. *Journal of Environmental Management*, 318: 115607.
- Limaki M.K., Nimvari M.E.H., Alavi S.J., Mataji A. & Kazemnezhad F., 2021 – Potential elevation shift of oriental beech (*Fagus orientalis* L.) in Hyrcanian mixed forest ecoregion under future global warming. *Ecological Modelling*, 455: 109637.
- Lippi C.A., Stewart-Ibarra A.M., Loo M.E.F.B., Zambrano J.E.D., Lopez N.A.E., Blackburn J.K. & Ryan S.J., 2019 – Geographic shifts in aedes aegypti habitat suitability in Ecuador using larval surveillance data and ecological niche modeling: implications of climate change for public health vector control. *PLoS Neglected Tropical Diseases*, 13 (4): e0007322.
- Littlefield C.E., Krosby M., Michalak J.L. & Lawler J.J., 2019 – Connectivity for species on the move: supporting climate-driven range shifts. *Frontiers in Ecology and the Environment*, 17 (5): 270-278.
- Liu C., Newell G. & White M., 2019 – The effect of sample size on the accuracy of species distribution models: considering both presences and pseudo-absences or background sites. *Ecography*, 42 (3): 535-548.
- Liu J., Wilson M., Hu G., Liu J., Wu J. & Yu M., 2018 – How does habitat fragmentation affect the biodiversity and ecosystem functioning relationship? *Landscape Ecology*, 33: 341-352.
- Maassoumi A. A. and Ashouri P., 2022 – The hotspots and conservation gaps of the mega genus *Astragalus* (Fabaceae) in the old-world. *Biodiversity and Conservation*, 31: 2119-2139.
- Mahmoodi S., Heydari M., Ahmadi K., Khwarahm N.R., Karami O., Almasieh K., Naderi B., Bernard P. & Mosavi A., 2022 – The current and future potential geographical distribution of *Nepeta Crispa* willd., an endemic, rare and threatened aromatic plant of Iran: Implications for Ecological Conservation and Restoration. *Ecological Indicators*, 137: 108752.
- Mahmoodi S., Ahmadi K., Heydari M., Karami O., Esmailzadeh O. & Heung B., 2023 – Elevational shift of endangered European yew under climate change in Hyrcanian mountain forests: rethinking conservation-restoration strategies and management. *Forest Ecology and Management*, 529: 120693.
- Marsico T.D., Krimmel E. R., Carter J.R., Gillespie E.L., Lowe P.D., McCauley R., Morris A.B., Nelson G., Smith M. & Soteropoulos D.L. & Monfils A.K., 2020 – Small herbaria contribute unique biogeographic records to county, locality, and temporal scales. *American Journal of Botany*, 107 (11): 1577-1587.
- Mason R.E., Craine J.M., Lany N.K., Jonard M., Ollinger S.V., Groffman P.M., Fulweiler R.W., Angerer J., Read Q.D. & Reich P.B., Templer P.H. & Elmore A.J., 2022 – Evidence, causes, and consequences of declining nitrogen availability in terrestrial ecosystems. *Science*, 376: eabh3767.
- Masson-Delmotte V., Zhai P., Pirani A., Connors S.L., Péan C., Berger S., Caud N., Chen Y., Goldfarb L. & Gomis M., 2021 – Climate change 2021: the physical science basis. <<https://www.ipcc.ch/report/ar6/wg1/>>
- McCune J., 2016 – Species distribution models predict rare species occurrences despite significant effects of landscape context. *Journal of Applied Ecology*, 53 (6): 1871-1879.
- McPherson K., 2022 – Bringing home the lost vegetable sheep: a phylogenomic study of the Senecioneae genus *Haastia*. <<https://ir.canterbury.ac.nz/server/api/core/bitstreams/a4dbba05-e8c1-4921-ae5d-1c8d61879d73/content>>
- Mirhashemi H., Heydari M., Ahmadi K., Karami O., Kavgaci A., Matsui T. & Heung B., 2023 – Species distribution models of brant's oak (*Quercus Brantii* lindl.): the impact of spatial database on predicting the impacts of climate change. *Ecological Engineering*, 194: 107038.
- Molano-Flores B., Johnson S.A., Marcum P.B. & Feist M.A., 2023 – Utilizing herbarium specimens to assist with the listing of rare plants. *Frontiers in Conservation Science*, 4: 1144593.
- Naqinezhad A., De Lombaerde E., Gholizadeh H., Wasof S., Perring M. P., Meeussen C., De Frenne P. & Verheyen K., 2022 – The combined effects of climate and canopy cover changes on understorey plants of the hyrcanian forest biodiversity hotspot in northern Iran. *Global Change Biology*, 28 (3): 1103-1118.
- Nicholson E., Watermeyer K.E., Rowland J.A., Sato C.F., Stevenson S.L., Andrade A., Brooks T.M., Burgess N.D., Cheng S.T., Grantham H.S., Hill S.L., Keith D.A., Maron M., Metzke D., Murray N.J., Nelson C.R., Obura D., Plumptre A., Skowno A.K. & Watson J.E.M., 2021 – Scientific foundations for an ecosystem goal, milestones and indicators for the post-2020 global biodiversity framework. *Nature Ecology and Evolution*, 5: 1338-1349.
- Niknaddaf Z., Hemami M.R., Pourmanafi S. & Ahmadi M., 2023 – An integrative climate and land cover change detection unveils extensive range contraction in mountain newts. *Global Ecology and Conservation*, 48: e02739.
- Noori S., Hofmann A., Rödder D., Husemann M. & Rajaei H., 2024 – A window to the future: effects of climate change on the distribution patterns of Iranian Zygaenidae and their host plants. *Biodiversity and Conservation*, 33: 579-602.
- Nyumba T.O., Wilson K., Derrick C.J. & Mukherjee N., 2018 – The use of focus group discussion methodology: insights from two decades of application in conservation. *Methods in Ecology and Evolution*, 9 (1): 20-32.
- Ovaskainen O., Roy D.B., Fox R. & Anderson B.J., 2016 – Uncovering hidden spatial structure in species communities with spatially explicit joint species distribution models. *Methods in Ecology and Evolution*, 7 (4): 428-436.
- Oyster J.H., Keren I.N., Hansen S.J. & Harris R.B., 2018 – Hierarchical mark-recapture distance sampling to estimate moose abundance. *Journal of Wildlife Management*, 82 (8): 1668-1679.
- Palit K., Rath S., Chatterjee S. & Das S., 2022 – Microbial diversity and ecological interactions of microorganisms in the mangrove ecosystem: threats, vulnerability, and adaptations. *Environmental Science and Pollution Research*, 29: 32467-32512.
- Pesce S., Bérard A., Coutellec M., Hedde M., Langlais-Hesse A., Larras F., Leenhardt S., Mongruel R., Munaron D., Sabater S. & Gallai N., 2023 – Linking ecotoxicological effects on biodiversity and ecosystem functions to impairment of ecosystem services is a challenge: an illustration with the case of plant protection products. *Environmental Science and Pollution Research*, 32: 2773-2785.
- Petsch D.K., Cioneck V.M., Thomaz S.M. & dos Santos L., 2023 – Ecosystem services provided by river-floodplain ecosystems. *Hydrobiologia*, 850: 2563-2584.
- Plutino M., Bianchetto E., Durazzo A., Lucarini M., Lucini L. & Negri I., 2022 – Rethinking the connections between ecosystem services, pollinators, pollution, and health: focus on air pollution and its impacts. *International Journal of Environmental Research and Public Health*, 19 (5): 2997.
- Priscope N.G., Mapes K.L. & Woodward K.D., 2019 – Remote sensing of human-environment interactions in global change research: a review of advances, challenges and future directions. *Remote Sensing*, 11 (23): 2783.
- Qazi A. W., Saqib Z. & Zaman-ul-Haq M., 2022 – Trends in species distribution modelling in context of rare and endemic plants: a systematic review. *Ecological Processes*, 11: 40.

- Ribeiro B.R., Guidoni-Martins K., Tessarolo G., Velazco S.J.E., Jardim L., Bachman S.P. & Loyola R., 2022 – Issues with species occurrence data and their impact on extinction risk assessments. *Biological Conservation*, 273: 109674.
- Rocchini D., Petras V., Petrasova A., Horning N., Furtkevicova L., Neteler M., Leutner B. & Wegmann M., 2017 – Open data and open source for remote sensing training in ecology. *Ecological Informatics*, 40: 57-61.
- Rocha-Ortega M., Rodríguez P., Bried J., Abbott J. & Córdoba-Aguilar A., 2020 – Why do bugs perish? Range size and local vulnerability traits as surrogates of Odonata extinction risk. *Proceedings of the Royal Society B*, 287 (1924): 20192645.
- Rønsted N., Campbell R., DeMotta M., Edmonds M., Houck K., Kahukula Jr M., Mayfield K.K., Nyberg B., Oppenorth M., Walsh S.K., Wolkis D., Wood K.R. & Nagendra U., 2023 – Restoration of threatened plant species in Limahuli Valley on the Hawaiian I Sland of Kaua'i in the framework of the global tree assessment. *Plants, People, Planet*, 5 (4): 547-562.
- Rühm W., Friedl A.A. & Wojcik A., 2023 – Un sustainable development goals: establishment of an electronic 'collection' of papers published in radiation and environmental biophysics. *Radiation and Environmental Biophysics*, 62: 173-174.
- Safaei M., Rezayan H., Firouzabadi P.Z. & Sadidi J., 2021 – Optimization of species distribution models using a genetic algorithm for simulating climate change effects on Zagros forests in Iran. *Ecological Informatics*, 63: 101288.
- Sanguet A., Wyler N., Petitpierre B., Honeck E., Poussin C., Martin P. & Lehmann A., 2022 – Beyond topo-climatic predictors: does habitats distribution and remote sensing information improve predictions of species distribution models? *Global Ecology and Conservation*, 39: e02286.
- Saran S., Chaudhary S.K., Singh P., Tiwari A. & Kumar V., 2022 – A comprehensive review on biodiversity information portals. *Biodiversity and Conservation*, 31: 1445-1468.
- Shaikh S.F.E.A., See S.C., Richards D., Belcher R.N., Grêt-Regamey A., Torres MG. & Carrasco L.R., 2021 – Accounting for spatial autocorrelation is needed to avoid misidentifying trade-offs and bundles among ecosystem services. *Ecological Indicators*, 129: 107992.
- Shay J.E., Pennington L.K., Mandussi Montiel-Molina J.A., Toews D.J., Hendrickson B.T. & Sexton J.P., 2021 – Rules of plant species ranges: applications for conservation strategies. *Frontiers in Ecology and Evolution*, 9: 700962.
- Sillero N. & Barbosa A.M., 2021 – Common mistakes in ecological niche models. *International Journal of Geographical Information Science*, 35 (2): 213-226.
- Simião-Ferreira J., Nogueira D.S., Santos A.C., De Marco P. & Angelini R., 2018 – Multi-scale homogenization of caddisfly metacommunities in human-modified landscapes. *Environmental management*, 61: 687-699.
- Steinbauer M.J., Grytnes J., Jurasinski G., Kulonen A., Lenoir J., Pauli H., Rixen C., Winkler M., Bardy-Durchhalter M., Barni E., Bjorkman A.D., Breiner F.T., Burg S., Czortek P., Dawes M.A., Delimat A., Dullinger S., Erschbamer B., Felde V.A., Fernández-Arberas O., Fossheim K.F., Gómez-García D., Georges D., Grindrud E.T., Haider S., Haugum S.V., Henriksen H., Herreros M.J., Jaroszewicz B., Jaroszynska F., Kanka R., Kapfer J., Klanderud K., Kühn I., Lamprecht A., Matteodo M., Morra di Cella U., Normand S., Odland A., L Olsen S., Palacio S., Petey M., Piscová V., Sedlakova B., Steinbauer K., Stöckli V., Svenning J.C., Teppa G., Theurillat J.P., Vittoz P., Woodin S.J., Zimmermann N.E., Wipf S., 2018 – Accelerated Increase in Plant Species Richness on Mountain Summits Is Linked to Warming. *Nature*, 556: 231-234.
- Taleshi H., Jalali S.G., Alavi S.J., Hosseini S.M., Naimi B. & Zimmermann N.E., 2019 – Climate change impacts on the distribution and diversity of major tree species in the temperate forests of Northern Iran. *Regional Environmental Change*, 19: 2711-2728.
- Tanner E.P., Papeş M., Elmore R.D., Fuhlendorf S.D. & Davis C.A., 2017 – Incorporating abundance information and guiding variable selection for climate-based ensemble forecasting of species' Distributional Shifts. *PLoS One*, 12 (9): e0184316.
- Van Nuland M.E., Wooliver R.C., Pfennigwerth A.A., Read Q.D., Ware I.M., Mueller L., Fordyce J.A., Schweitzer J.A. & Bailey J.K., 2016 – Plant-soil feedbacks: connecting ecosystem ecology and evolution. *Functional Ecology*, 30 (7): 1032-1042.
- Veselova E. & Gaziulusoy I., 2021 – When a tree is also a multispecies collective, a photosynthesis process, and a carbon cycle: a systemic typology of natural nonhuman stakeholders when designing for sustainability. In: *Proceedings of Relating Systems Thinking and Design (RSD10)*. Diehl J. C., Tromp N. & Bijil-Brouwer v. d. (eds).
- Vilà-Cabrera A., Premoli A.C. & Jump A.S., 2019 – Refining predictions of population decline at species' rear edges. *Global Change Biology*, 25 (5): 1549-1560.
- Wang S., Loreau M., Arnoldi J., Fang J., Rahman K.A., Tao S. & de Mazancourt C., 2017 – An invariability-area relationship sheds new light on the spatial scaling of ecological stability. *Nature Communications*, 8: 15211.
- Wen Z., Yang Q., Quan Q., Xia L., Ge D. & Lv X., 2016 – Multiscale partitioning of small mammal  $\beta$ -diversity provides novel insights into the quaternary faunal history of Qinghai-Tibetan plateau and Hengduan mountains. *Journal of Biogeography*, 43 (7): 1412-1424.
- Yu D., Liu Y., Shi P. & Wu J., 2019 – Projecting impacts of climate change on global terrestrial ecoregions. *Ecological Indicators*, 103: 114-123.
- Zeraatkar A. & Khajoei Nasab F., 2023 – Mapping the habitat suitability of endemic and sub-endemic almond species in Iran under current and future climate conditions. *Environment, Development and Sustainability*, 26: 14859-14876.
- Zhong Y., Xue Z., Jiang M., Liu B. & Wang G., 2021 – The application of species distribution modeling in wetland restoration: a case study in the songnen plain, northeast China. *Ecological Indicators*, 121: 107137.