



Natural History Sciences

<https://sisn.pagepress.org/nhs>

eISSN 2385-0922

Publisher's Disclaimer. E-publishing ahead of print is increasingly important for the rapid dissemination of science. The Early Access service lets users access peer-reviewed articles well before print/regular issue publication, significantly reducing the time it takes for critical findings to reach the research community.

These articles are searchable and citable by their DOI (Digital Object Identifier).

Natural History Sciences is, therefore, E-publishing PDF files of an early version of manuscripts that have undergone a regular peer review and have been accepted for publication, but have not been through the copyediting, typesetting, pagination, and proofreading processes, which may lead to differences between this version and the final one.

The final version of the manuscript will then appear in a regular issue of the journal.

The E-publishing of this PDF file has been approved by the authors.

Natural History Sciences 2026 [Online ahead of print]

Please cite this article as:

Filippo Scafidi, Alfonso La Rosa, Salvatore Pasta. **Typification of the name *Herniaria empedocleana* (Caryophyllaceae).** *Natural History Sciences* doi: 10.4081/nhs.2026.996

Submitted: 22 January 2026

Accepted: 29 June 2026

 © the Author(s), 2026
Licensee PAGEPress, Italy

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Typification of the name *Herniaria empedocleana* (Caryophyllaceae)

Filippo Scafidi^{1*}, Alfonso La Rosa², Salvatore Pasta³

¹ Orto Botanico di Palermo, UNIPA Heritage, Sistema Museale di Ateneo, Via Lincoln 2, I-90133 Palermo, Italia.

² Dipartimento di Scienze e Tecnologie Biologiche Chimiche e Farmaceutiche (STEBICEF), Università di Palermo, Via Archirafi 38, I-90123 Palermo, Italia.

³ Consiglio Nazionale delle Ricerche (CNR), Istituto di Bioscienze e BioRisorse (IBBR), Sede di Palermo, Via Ugo La Malfa 153, I-90146 Palermo, Italia.

* Corresponding author: filippo.scafidi@unipa.it

© 2026 Filippo Scafidi, Alfonso La Rosa, Salvatore Pasta

Abstract - *Herniaria fontanesii* subsp. *empedocleana* is a Sicilian endemic vascular plant which has probably gone extinct during the last decades. In this contribution we propose a lectotype for the basionym, *Herniaria empedocleana* described by Lojacono-Pojero, whose selected specimen is kept in PAL. Some details are also provided on the figure and scientific activity of Lojacono-Pojero, one of the most influential Sicilian botanists of the second half of the 19th century. Additionally, some considerations are shared on the need to translate the work of typification and identification of *loci classici* into an effective tool for safeguarding the naturalistic values of the territory.

Keywords: herbaria, history of botany, nature conservation, Sicily, typification.

Riassunto - Tipificazione del nome *Herniaria empedocleana* (Caryophyllaceae).

Tipificazione del nome *Herniaria empedocleana*, basionimo di *Herniaria fontanesii* subsp. *empedocleana* (Caryophyllaceae), una pianta probabilmente estinta un tempo endemica della Sicilia. *Herniaria fontanesii* subsp. *empedocleana* è una pianta vascolare endemica siciliana, probabilmente estintasi negli ultimi decenni. In questo contributo proponiamo un lectotipo per il basionimo, *Herniaria empedocleana* descritto da Lojacono-Pojero, il cui esemplare selezionato è conservato in PAL. Vengono inoltre forniti alcuni dettagli sulla figura e sull'attività scientifica di Lojacono-Pojero, uno dei botanici siciliani più influenti della seconda metà del XIX secolo. Inoltre, vengono condivise alcune considerazioni sulla necessità di tradurre il lavoro di tipificazione e identificazione dei *loci classici* in uno strumento efficace per la salvaguardia dei valori naturalistici del territorio.

Parole chiave: conservazione della natura, erbari, Sicilia, storia della botanica, tipificazione.

INTRODUCTION

Within the Mediterranean Basin biodiversity hotspot (Myers *et al.*, 2000), the Italian peninsula is one of the most important centers of diversity and floristic richness (D'Antraccoli *et al.*, 2024). Furthermore, the presence of several key areas of plant diversity, which serve as both refuges and crossroads for plant dispersal, promotes active plant speciation (Médail & Quézel, 1997), making it still possible to describe taxa that are new to science (e.g. Bartolucci *et al.*, 2021, 2023; Brullo *et al.*, 2023; Cambria *et al.*, 2024, 2025; Bartolucci & Conti, 2025).

According to the most recent assessment, around 1,700 species and subspecies are endemic to Italy (Bartolucci *et al.*, 2024), 13 of which have been reported to be extinct in nature (Orsenigo *et al.*, 2018; Bartolucci *et al.*, 2024). One of them is *Herniaria fontanesii* Gay subsp. *empedocleana* (Lojac.) Brullo, last observed in 1980s () but not found in the last years despite repeated searches in the field

(S. Brullo and S. Cambria, *pers. comm.*). According to POWO (2017+), this subspecies belongs to the *H. fontanesii* aggregate that includes three taxa: *H. fontanesii* subsp. *fontanesii*, occurring in NW Africa and Canary Islands, *H. fontanesii* subsp. *almeriana* Brummitt & Heywood, growing in SE Spain and N Morocco, and *H. fontanesii* subsp. *empedocleana*, endemic to Sicily (southern Italy). A description of the main morphological traits of *H. fontanesii* subsp. *empedocleana* is available in Brullo & Brullo (2020).

Herniaria fontanesii subsp. *empedocleana* used to grow on marly and clayey soils from 0 to 20 m a.s.l. as a component of a nitro-xerophilous shrubland community dominated by woody chenopods corresponding to the association *Salsolo oppositifoliae-Limonietum opulenti* Brullo, Grillo & Scalia 1980 (Brullo *et al.*, 1980). This plant was first described as *Herniaria empedocleana* Lojac. by Lojacono-Pojero (1884: 284). Twenty years later, Lojacono-Pojero himself (1907: 264) reported this plant to grow only on the arid marls of one single locality, known as “Salatore”, located close to the harbour of Porto Empedocle near Agrigento (southern Sicily).

As part of an ongoing process of typification of the names referring to Sicilian plants published by Lojacono-Pojero (e.g., Domina & Mazzola, 2007; Aghababian *et al.*, 2008; Domina *et al.*, 2014a, 2014b, 2016, 2017; Di Gristina *et al.*, 2017; Scafidi *et al.*, 2020; Iamónico & Domina, 2023; Romero Zarco, 2023; Pasta & Troia, 2024, and references therein), the selection of the lectotype of the name *H. empedocleana*, currently not designated (Peruzzi *et al.*, 2015), is here done and discussed.

MATERIAL AND METHODS

This work is based on the analysis of Lojacono Pojero’s protologue and the examination of relevant literature. In order to trace the original material of *Herniaria empedocleana*, we performed a survey in PAL (codes follow Thiers, 1997+), where Lojacono-Pojero worked and where several specimens collected by him are kept (Stafleu & Cowan, 1981). Furthermore, we also checked the presence of specimens at B, BM, DS, E, FI, G, GE, K, L, MANCH, P, PI, NAP, W, and WU, where several Sicilian duplicates are kept, in order to trace further original material. The typification procedure follows the Madrid Code (Turland *et al.*, 2025), hereinafter ICN (International Code of Nomenclature).

RESULTS

In his protologue, Lojacono-Pojero (1884) provided detailed diagnosis, followed by the locus classicus “In aridis marnaceis mari finitimis cum Reaumuria vermiculata ac Statice Smithii, prope Porto Empedocle. Junio 1883”. Furthermore, the author explains in detail the main morphological differences between *H. empedocleana* and *H. fontanesii* Gay.

The herbarium search allowed us to trace one well preserved sheet stored at PAL (barcode PAL47570). No further original material has been found in the other investigated herbaria. The sheet from PAL contains various specimens mounted together, showing every organ of the plant except for the roots. The label hosts two different handwritings: the first is by G. Bonafede, gardener at the Botanical Garden of Palermo under the direction of Agostino Todaro: “Herniaria. Porto Empedocle/Bonafede. Aprile 1882”. Further information was added afterwards by d Lojacono-Pojero, whose handwriting is unmistakable: after the name of the genus *Herniaria* he added “*empedocleana* Lojac. Nat[.] Sicil[.]”, providing further details on the exact location of the plant, found and collected by Bonafede “[in] mezzo al Salatore Lojac.!” [= in the middle of the Salatore, personally seen by Lojacono-Pojero himself].

DISCUSSION

The explicit mention – made on the label by Lojacono-Pojero himself – of ‘Nat Sic.’ clearly refers to the paper, published in the journal ‘Il Naturalista siciliano’ where he first described *H. empedocleana*. Hence, the material fixed on this sheet is eligible for typification purposes. Moreover, PAL47570

perfectly matches the protologue and corresponds to the current application of the name (see e.g., Brullo *et al.*, 1980; Pignatti *et al.*, 2017; Brullo & Brullo, 2020) and therefore is here selected as the lectotype according to Art. 9.3 of the ICN.

Herniaria empedocleana Lojac., *Naturalista sicil.* 3(10): 284. 1884.

≡ *Herniaria fontanesii* subsp. *empedocleana* (Lojac.) Brullo, *Giorn. Bot. Ital.* n.s. 114(1-2): 46. 1980.

Lectotype (designated here): [Italy, Sicily] “*Herniaria* [manu Bonafede] *empedocleana* Lojac, Nat[.] Sicil[.] [manu Lojacono-Pojero] Porto Empedocle, *Bonafede*. Aprile 1882. Mezzo i Salvatore Lojac. [manu Lojacono-Pojero] (PAL No. PAL47570!, larger upper right fragment) (Fig. 1) (image available at https://herbarium.unipa.it/herbarium_vdetails_en2.asp?idmode=adv&id=58631).

Michele Lojacono Pojero (Palermo 1853-Messina 1919) has been one of the most influential Sicilian botanists of the second half of the 19th century. As a pupil of Agostino Todaro, director of the Royal Botanical Garden of Palermo between 1856 and 1892, he started investigating the Sicilian flora (not only vascular plants but also mosses and liverworts) around 1877. In 1883 he obtained a lectureship in botany (Dia, 1987), without ever receiving a salary. For most of his life he lived in Palermo, where he drew most of the economic resources necessary to finance his prolific research (Domina *et al.*, 2014b) from the trade of dry specimens of Sicilian plants sold to herbaria throughout Europe and even overseas. The disagreements and misunderstandings with his old mentor became even more bitter in 1892 with Todaro’s successor, Antonino Borzi (Domina *et al.*, 2014a). This situation prompted him to move to Messina in 1913, where he obtained a position as teacher of Natural Sciences at the Regio Istituto Tecnico (= Royal Technical Institute) and spent the last years of his life.

Apart from a few collections made in his youth, such as the one made in April 1877 on the Aeolian Islands, most of Lojacono-Pojero’s dry specimens are now preserved in European and even extra-European herbaria to which he sold his collections. He was probably forced to do so as a means of self-financement, having failed to obtain a paid position at the Botanical Garden due to Todaro’s obstructionism first and to Borzi’s subsequent hostility. It is therefore not surprising that PAL harbors very few types of the hundreds of taxa (species, varieties, forms) described by Lojacono-Pojero. The presence of original material referred to *Herniaria empedocleana* in PAL at first glimpse may look like an exception, but in fact it confirms this rule, as the specimen designated as lectotype comes from other people’s collections.

CONCLUSIONS

Sicily is not only one of the botanically richest regions in Italy, but also one with the highest number of extinct species or species that have not been observed for a long time (Domina *et al.*, 2025). In this regard, it should be noted that the probable extinction of *Herniaria fontanesii* subsp. *empedocleana* is not due to natural processes but is the result of the physical destruction of the territory where the only known population occurred. Between the late 19th century and the 1960s, in fact, the natural environment of the Porto Empedocle area was literally wiped out to make way for industrial buildings devoted to sulphur refining (Syed *et al.*, 2024). The rapid but short-lived industrial development of the area in turn led to chaotic urban sprawl and infrastructure development, giving the coup de grace to the local landscape and triggering the disappearance or irreversible degradation of most of the local ecosystems.

Over the last ten years, the Italian botanical community has made a considerable effort to highlight the importance of typification work and the identification of the *loci classici* of vascular plants endemic to Italy (Peruzzi *et al.*, 2015) or described from Italy (Peruzzi *et al.*, 2019). The richness of species – and of endemic ones in particular – in our flora should not just be something to boast about triumphantly at specialist conferences. Instead, it should remind not only botanists, but also politicians, local administrators, stakeholders and citizens of their duty to protect these places. The

importance of *loci classici* becomes crucial in the case of endemic species that count only one or a few populations, whose protection is our full and sole responsibility (Brundu *et al.*, 2017).

In recent decades, several species new to science have been described whose past presence is now documented only by herbarium material, while these plants are no longer present in nature, having become extinct globally as a result of the destruction or profound alteration of the territories where they used to live: this is the case, for example, of *Limonium catanense* (Tineo ex Lojac.) Brullo, which probably disappeared at the beginning of the 20th century from the coastal area of the port of Catania, or *Suaeda kocheri* Guss. ex C.Brullo, Brullo & Giusso, once present near the ports of Lampedusa and Porto Empedocle, where it seems to have disappeared definitively over the last few decades (Brullo *et al.*, 2011). Identifying the locations where extremely rare plants have been collected and from where they have been described should be the first step in adopting effective strategies to safeguard or increase their integrity, allowing these rare plants to grow in habitats that guarantee their growth, reproduction and free dispersal. Otherwise, scientific research applied to the nomenclature and classification of plants risks remaining a sterile academic exercise with no practical impact, and type labels may sound like sad epitaphs.

Acknowledgements

The authors are grateful to Alfredo Carratello, curator of the *Herbarium Mediterraneum Panormitanum* (PAL) for his support during the study of the *exsiccata*. They also wish to thank Salvatore Brullo and Salvatore Cambria (University of Catania) for sharing their personal information on this plant.

REFERENCES

- Aghababian M., Greuter W., Mazzola P. & Raimondo F.M., 2008 – Typification of names of Compositae taxa described from Sicily by Michele Lojacono-Pojero. *Flora Mediterranea*, 18: 513-528.
- Bartolucci F. & Conti F., 2025 – *Thymus apulus* (T. sect. *Hyphodromi*, Lamiaceae), a new species from southern Italy. *Plants*, 14 (23), 3584. <<https://doi.org/10.3390/plants14233584>>
- Bartolucci F., Galasso G., Peruzzi L. & Conti F., 2021 – Report 2020 on plant biodiversity in Italy: native and alien vascular flora. *Natural History Sciences*, 8 (1): 41-54. <<https://doi.org/10.4081/nhs.2021.520>>
- Bartolucci F., Galasso G., Peruzzi L. & Conti F., 2023 – Report 2021 on plant biodiversity in Italy: native and alien vascular flora. *Natural History Sciences*, 10 (1) (2022): 41-50. <<https://doi.org/10.4081/nhs.2022.623>>
- Bartolucci F., Peruzzi L., Galasso G., Alessandrini A., Ardenghi N.M.G., Bacchetta G., Banfi E., Barberis G., Bernardo L., Bouvet D., Bovio M., Calvia G., Castello M., Cecchi L., Del Guacchio E., Domina G., Fascetti S., Gallo L., Gottschlich G., Guarino R., Gubellini L., Hofmann N., Iberite M., Jiménez-Mejías P., Longo D., Marchetti D., Martini F., Masin R.R., Medagli P., Peccenini S., Prosser F., Roma-Marzio F., Rosati L., Santangelo A., Scoppola A., Selvaggi A., Selvi F., Soldano A., Stinca A., Wagensommer R.P., Wilhalm T. & Conti F., 2024. – A second update to the checklist of the vascular flora native to Italy. *Plant Biosystems*, 158 (2): 219-296. <<https://doi.org/10.1080/11263504.2024.2320126>>
- Brullo C. & Brullo S., 2020 – Flora endemica illustrata della Sicilia. *Laruffa Editore*, Reggio Calabria.
- Brullo S., Grillo M. & Scalia C., 1980 – Su una nuova associazione dei *Pegano-Salsoletea* osservata presso Porto Empedocle (Agrigento). *Giornale Botanico Italiano*, 114 (1-2): 43-52. <<https://doi.org/10.1080/11263508009426433>>
- Brullo C., Brullo S. & Giusso del Galdo G., 2011 – *Suaeda kocheri* (Chenopodiaceae), a new species from Sicily. *Phyton*, 51 (1): 89-94.
- Brullo S., Brullo C., Cambria S., Tomaselli V., Crisafulli A., Siracusa G., Minissale P. & Giusso del Galdo G., 2023 – Taxonomic and ecological remarks on *Solenopsis bivonae* species complex

- (Campanulaceae). *PhytoKeys*, 229: 77-111. <<https://doi.org/10.3897/phytokeys.229.104324>>
- Brundu G., Peruzzi L., Domina G., Bartolucci F., Galasso G., Peccenini S., Raimondo F.M., Albano A., Alessandrini A., Banfi E., Barberis G., Bernardo L., Bovio M., Brullo S., Brunu A., Camarda I., Carta L., Conti F., Croce A., Iamónico D., Iberite M., Iiriti G., Longo D., Marsili S., Medagli P., Mariotti M. G., Pennesi R., Pistarino A., Salmeri C., Santangelo A., Scassellati E., Selvi F., Stinca A., Vacca G., Villani M., Wagensommer R.P. & Passalacqua N.G., 2017 – At the intersection of cultural and natural heritage: Distribution and conservation of the type localities of Italian endemic vascular plants. *Biological Conservation*, 214: 109-118. <<https://doi.org/10.1016/j.biocon.2017.07.024>>
- Cambria S., Giusso del Galdo G., Minissale P., Tavilla G. & Salmeri C., 2024 – *Cyclamen brulloi* (Primulaceae), a new species from Sicily (Italy). *Kew Bulletin*, 79 (3): 653-670. <<https://doi.org/10.1007/s12225-024-10207-2>>
- Cambria S., Ranno V., Minissale P., Giusso del Galdo G., Salmeri C., Monari D., La Rosa A., Siracusa G. & Scafidi F., 2025 – Notes on the *Vicia tenuifolia* complex with the description of a new species, *V. barbatoï* from Sicily. *Taxonomy*, 5 (4), 63. <<https://doi.org/10.3390/taxonomy5040063>>
- D'Antraccoli M., Peruzzi L., Conti F., Galasso G., Roma-Marzio F. & Bartolucci F., 2024 – Floristic richness in a Mediterranean hotspot: a journey across Italy. *Plants*, 13 (1), 12. <<https://doi.org/10.3390/plants13010012>>
- Dia M.G., 1987 – L'opera di Michele Lojacono-Pojero per la illustrazione della Flora Sicula. In: I naturalisti e la cultura scientifica nell'Ottocento. Liotta G. (ed.). *Atti del convegno*, Palermo: 495-503.
- Di Gristina E., Domina G., Raimondo F.M. & Scafidi F., 2017 – Typification of the name *Abies nebrodensis* (Pinaceae). *Taxon*, 66 (4): 970-972. <<https://doi.org/10.12705/664.11>>
- Domina G. & Mazzola P., 2007 – The genus *Orobanche* in Sicily. Taxa described by V. Tineo and M. Lojacono Pojero. *Boccone*, 21: 223-232.
- Domina G., Greuter W., Mazzola P. & Raimondo F.M., 2014a – Names of Italian vascular plants published by Michele Lojacono Pojero. *Flora Mediterranea*, 24: 215-232.
- Domina G., Greuter W., Mazzola P. & Raimondo F.M., 2014b – La ricerca dei tipi di Michele Lojacono Pojero nell'“Herbarium Mediterraneum Panormitanum”. *Museologia Scientifica Memorie*, 11: 81-84.
- Domina G., Greuter W. & Raimondo F.M., 2016 – Types of names of taxa belonging to the *Centaurea cineraria* group (*Compositae*) described from Sicily. *Willdenowia*, 46 (1): 23-26. <<https://doi.org/10.3372/wi.46.46102>>
- Domina G., Scibetta S., Scafidi F. & Giovino A., 2017 – Contribution to the identification of *Dianthus rupicola* (Caryophyllaceae) subspecies using morphological and molecular approaches. *Phytotaxa*, 291 (1): 17-32. <<https://doi.org/10.11646/phytotaxa.291.1.2>>
- Domina G., Abeli T., Alessandrini A., Angiolini C., Argenti C., Astuti G., Bacchetta G., Bagella S., Ballelli S., Banfi E., Barberis G., Barone G., Bernardo L., Bertolli A., Bianchi E., Bonari G., Bouvet D., Bovio M., Briozzo I., Brullo C., Brullo S., Calvia G., Caria M.C., Caputo P., Caruso G., Casazza G., Cecchi L., Cibeï C., Cogoni D., Conti F., Crisafulli A., Croce A., Dagnino D., Del Guacchio E., Di Gristina E., Di Marzio P., Fanfarillo E., Farris E., Fascetti S., Fenu G., Ferrando U., Ferretti G., Fiaschi T., Fortini P., Fois M., Franzoni J., Galasso G., Galesi R., Ganz C., Gargano D., Giacò A., Gianguzzi L., Giardini M., Giusso del Galdo G., Guarino R., Gubellini L., Hofmann N., Iamónico D., Iberite M., Laface V.L.A., Lallai A., Lastrucci L., Lavezzo P., Lazzaro L., Longo D., Lucchese F., Maiorca G., Manca M., Marchetti D., Marsili S., Martini F., Mascia F., Medagli P., Mondello F., Montagnani C., Mugnai M., Musarella C.M., Nicoletta G., Orsenigo S., Passalacqua N.G., Peccenini S., Pennesi R., Peruzzi L., Picone R.M., Pinzani L., Podda L., Prosser F., Riviaccio G., Roma-Marzio F., Rosati L., Salerno G., Santangelo A., Sciandrello S., Selvaggi A., Selvi F., Spampinato G., Stinca A., Tardella F.M., Tavilla G., Tilia A., Tondi G., Tognon G., Urbani M., Villani M., Wilhalm T., Zappa E. & Bartolucci F., 2025 – Refining the inventory of the vascular flora of Italy: Unconfirmed,

- doubtful, excluded or extinct taxa. *Phytotaxa*, 723 (1).
<<https://doi.org/10.11646/phytotaxa.723.1.1>>
- Iamonico D. & Domina G., 2023 – Studies on the genus *Atriplex* (Chenopodiaceae) in Italy. VIII. Names published by Vincenzo Tineo and Michele Lojacono-Pojero. *Flora Mediterranea*, 33: 49-59. <<https://doi.org/10.7320/FIMedit33.049>>
- Lojacono-Pojero M., 1884 – Studi su piante critiche rare o nuove della flora di Sicilia. *Il Naturalista Siciliano*, 3 (10): 282-286.
- Lojacono-Pojero M., 1907 – *Herniaria* Lin. In: Flora sicula o descrizione delle piante spontanee o indigenate in Sicilia. Lojacono-Pojero M. (ed.). *Tipo-Litografia Salvatore Bizzarrilli*, Palermo, 2 (2): 261-264.
- Médail F. & Quézel P., 1997 – Hot-spots analysis for conservation of plant biodiversity in the Mediterranean Basin. *Annals of the Missouri Botanical Garden*, 84 (1): 112-127. <<https://doi.org/10.2307/2399957>>
- Myers N., Mittermeier R.A., Mittermeier C.G., Da Fonseca G.A.B. & Kent J., 2000 – Biodiversity hotspots for conservation priorities. *Nature*, 403: 853-858. <<https://doi.org/10.1038/35002501>>
- Orsenigo S., Montagnani C., Fenu G., Gargano D., Peruzzi L., Abeli T., Alessandrini A., Bacchetta G., Bartolucci F., Bovio M., Brullo C., Brullo S., Carta A., Castello M., Cogoni D., Conti F., Domina G., Foggi B., Gennai M., Gigante D., Iberite M., Lasen C., Magrini S., Perrino E.V., Prosser F., Santangelo A., Selvaggi A., Stinca A., Vagge I., Villani M., Wagensommer R.P., Wilhalm T., Tartaglini N., Duprè E., Blasi C. & Rossi G., 2018 – Red Listing plants under full national responsibility: extinction risk and threats in the vascular flora endemic to Italy. *Biological Conservation*, 224: 213-222. <<https://doi.org/10.1016/j.biocon.2018.05.030>>
- Pasta S. & Troia A., 2024 – Typification of the name *Cistus ×skanbergii* Lojac., a rare rockrose extinct in its type locality. *Natural History Sciences*, 11 (1): 3-8. <<https://doi.org/https://doi.org/10.4081/nhs.2024.665>>
- Peruzzi L., Domina G., Bartolucci F., Galasso G., Peccenini S., Raimondo F.M., Albano A., Alessandrini A., Banfi E., Barberis G., Bernardo L., Bovio M., Brullo S., Brundu G., Brunu A., Camarda I., Carta L., Conti F., Croce A., Iamonico D., Iberite M., Iiriti G., Longo D., Marsili S., Medagli P., Pistarino A., Salmeri C., Santangelo A., Scassellati E., Selvi F., Soldano A., Stinca A., Villani M.C., Wagensommer R.P. & Passalacqua N.G., 2015 – An inventory of the names of vascular plants endemic to Italy, their loci classici and types. *Phytotaxa*, 196 (1): 1-217. <<https://doi.org/10.11646/phytotaxa.196.1.1>>
- Peruzzi L., Galasso G., Domina G., Bartolucci F., Santangelo A., Alessandrini A., Astuti G., D'antraccoli M., Roma-Marzio F., Ardenghi N.M.G., Barberis G., Conti F., Bernardo L., Peccenini S., Stinca A., Wagensommer R.P., Bonari G., Iamonico D., Iberite M., Viciani D., Del Guacchio E., Giusso Del Galdo G., Lastrucci L., Villani M., Brunu A., Magrini S., Pistarino A., Brullo S., Salmeri C., Brundu G., Clementi M., Carli E., Vacca G., Marcucci R., Banfi E., Longo D., Di Pietro R. & Passalacqua N.G., 2019 – An inventory of the names of native, non-endemic vascular plants described from Italy, their loci classici and types. *Phytotaxa*, 410 (1): 1-215. <<https://doi.org/10.11646/phytotaxa.410.1.1>>
- Pignatti S., Guarino R. & La Rosa M., 2017 – Flora d'Italia. Volume 2. *Edagricole*, Bologna, 2^a edizione.
- POWO, 2017+ – Plants of the world online. *Board of Trustees of the Royal Botanic Gardens, Kew*. <<https://powo.science.kew.org/>> (Retrieved in January 3, 2026).
- Romero Zarco C., 2023 – *Bromus cincinnatus* (Poaceae): perennial oat-grass or annual brome-grass? *Mediterranean Botany*, 44, e84748. <<https://doi.org/10.5209/mbot.84748>>
- Scafidi F., Ferraro V. & Di Gristina E., 2020 – Typification of the name *Eryngium siculum* (Apiaceae). *Phytotaxa*, 461 (3): 222-224. <<https://doi.org/10.11646/phytotaxa.461.3.10>>
- Stafleu F.A. & Cowan R.S., 1981 – Taxonomic literature. A selective guide to botanical publications and collections with dates, commentaries and types. Vol. III Lh-O (second edition). *Bohn, Scheltema & Holkema*, Utrecht.
- Syed K., Andrade M.J. & Casares L.M., 2024 – Dismissed industrial areas: the urban project of the

Montedison factory of Akragas in Porto Empedocle between the past, present, and future. *Buildings*, 14 (12), 3895. <<https://doi.org/10.3390/buildings14123895>>

Thiers B., 1997+ – *Index herbariorum*: a global directory of public herbaria and associated staff. *New York Botanical Garden*. <<http://sweetgum.nybg.org/science/ih/>> (Retrieved in January 3, 2026).

Turland N.J., Wiersema J.H., Barrie F.R., Gandhi K.N., Gravendyck J., Greuter W., Hawksworth D.L., Herendeen P.S., Klopper R.R., Knapp S., Kusber W.-H., Li D.-Z., May T.W., Monro A.M., Prado J., Price M.J., Smith G.F. & Zamora Señoret J.C. (eds.), 2025 – International Code of Nomenclature for algae, fungi, and plants (Madrid Code). *University of Chicago Press, Regnum Vegetabile*, 162. <<https://doi.org/10.7208/chicago/9780226839479.001.0001>>




Fig. 1 - Sheet bearing the lectotype (larger upper right fragment) of the name *Herniaria empedocleana* (PAL47570, published with permission). / Foglio recante il lectotipo (frammento superiore destro più grande) del nome *Herniaria empedocleana* (PAL47570, pubblicato con autorizzazione).